## **Ustekinumab (Stelara)**

Provider Order Form rev. 01/02/2024

Provider Name (print)

PATIENT INFORMATION	Referral Status	: □ New Re	eferral 🗆 Upda	ated Order	☐ Order Renewal
Patient Name:		DOB:	Pat	tient Phone:	
Patient Address:	Patient Email:				
Allergies:		□ NKDA	Weight (lbs/kg)	: He	eight (in/cm):
Sex: $\square$ M / $\square$ F Date of Last Infusion:	Next Due Dat	e:	Preferred Loc		
DIAGNOSIS (Please provide ICD-10 code in space		C I''			
Crohn's Disease:	Ulcerative Colitis:				
Plaque Psoriasis:	Psoriati	c Arthritis:			_
REQUIRED INFORMATION (Choose one)  □ Patient will self-administer subcutaneous medication (Referring provider will coordinate with specialty pharmacy) □ Patient would like in-office injection medication (NOTE: some insurance providers may require attestation from provider stating		IV DOSE PRE-MEDICATION ORDERS  ☐ Tylenol ☐ 500mg / ☐ 650mg PO ☐ Loratadine 10mg PO ☐ Pepcid 20mg ☐ PO / ☐ IVP ☐ Benadryl ☐ 25mg / ☐ 50mg ☐ PO / ☐ IVP			
patient cannot self-administer with reason why such as needle		☐ Solumedrol ☐ 40mg / ☐ 125mg IVP			
phobia or low dexterity.)		☐ Other:			
THERAPY ADMINISTRATION & DOSING  For Crohn's/Ulcerative Colitis:  ☐ Induction: Administer Stelara mixed in 250ml 0.9% NS over 1 hour on week 0, one time dose only:  ☐ 260mg IV x1 dose (weight of up to 55kg) ☐ 390mg IV x1 dose (weight of 55kg to 85kg) ☐ 520mg IV x1 dose (weight greater than 85kg) ☐ Maintenance: Administer Stelara 90mg subcutaneously every 8 weeks		NURSING  ☑ Hold infusion and notify provider for:  • Signs/symptoms of illness/active infection or cough, night sweats, unexplained weight loss  • Planned/recent surgical procedures, recent live vaccinations, or neurological changes  ☑ Provide nursing care per Nursing Procedure, including Hypersensitivity Reaction Management Protocol and post-procedure observation			
		For Plaque Psoriasis/Psoriatic Arthritis:  Induction: Administer Stelara subcutaneously on we week 4:  Induction: Administer Stelara subcutaneously on we week 4:  Induction: Administer Stelara Stelara 100 weight less than 100 weeks (weight less than 100kg)  Induction: Administer Stelara 45mg subcutaneously (weight less than 100kg)  Induction: Administer Stelara 90mg subcutaneoweeks (weight greater than 100kg)	00kg) n 100kg) usly every 12		
PROVIDER INFORMATION		Du-f	Contact Fun		
Preferred Contact Name: Ordering Provider:		Preferred Contact Email: Provider NPI:			
Referring Practice Name:	Dł	Phone:		Fax:	
Practice Address:		ty:	State:		Zip Code:
					•
REQUIRED DOCUMENTATION CHECKLIST (A Required Documentation: Patient demos, copy of for treatment failures or contraindications, Colonoscop Required Labs: TB, Hep B ESR, CRP, for RA: RF, CCP,	ront and back of p y, reason patient	orimary and s is unable to s	econdary insurar	nce, 2 most re	ecent OVN including

Date

**Provider Signature**